

RESEARCH LETTER

Modeling of a Novel Patient-Based *MYO5B* Point Mutation Reveals Insights Into MVID Pathogenesis



Microvillus inclusion disease (MVID) is a congenital disorder that presents with severe secretory diarrhea, typically within the first few hours of life. Inactivating mutations of myosin VB (*MYO5B*) cause MVID, which usually requires lifetime total parenteral nutrition or small-bowel transplantation.¹ A recent analysis of 114 *MYO5B* mutations identified that most mutations were compound heterozygous.² Although previously developed homozygous *MYO5B* knockout animal models showed success in recapitulating MVID,³ there is a lack of research into the phenotypes caused by compound heterozygous mutations of *MYO5B*.

Targeted exome sequencing in an MVID patient and both parents identified 2 variants in the *MYO5B* gene: c.1821delG (p.S608HfsX14), likely a de novo mutation that leads to early truncation, and c.1555G>A (p.G519R) (Supplementary Figure 1A), which was maternally inherited. This patient mutation is referred to as MYO5B(G519R) in this study. The patient had copious diarrhea from birth with non-ion gap acidosis and intermittent feeding intolerance. An esophagogastroduodenoscopy and flexible sigmoidoscopy at 3 weeks of age were grossly normal. Biopsy specimens showed villus blunting and abnormal accumulation of periodic acid-Schiff (PAS)-positive vesicles in the enterocytes (Figure 1A). Transmission electron microscopy showed disorganized and shortened microvilli, an abnormal accumulation of subapical vesicles, and multivesicular bodies. Brush-border dipeptidyl peptidase 4 (DPP4) and terminal web α -actinin-4

were obscure in the MVID patient tissues (Figure 1B). Intriguingly, A-kinase anchoring protein (AKAP) 350 expression was found around the inclusion-like structure, suggesting that the A-kinase anchoring protein-mediated scaffolding pathway is involved in microvillus inclusion formations.

Multiplexed immunofluorescence staining (MxIF) showed MVID characteristics on a single slide of duodenum biopsy specimens using 15 markers, including previously known and unexplored proteins in MVID tissues (Figure 1C and D). Basolateral nutrient transporters other than Na-K-adenosine triphosphatase have not been characterized in MVID tissues.⁴ The present MxIF demonstrates a remarkable decrease in GLUT2-expressing enterocytes in the MVID patient with MYO5B(G519R), implicating a deficit in enterocyte maturation that could contribute to malabsorption.^{5,6} Consistent with reports from MVID patient tissues with other *MYO5B* mutations,^{5,6} apical nutrient transporters, such as sodium-dependent glucose transporter 1 (SGLT1) and sodium hydrogen antiporter 3 (NHE3), were internalized, and brush-border markers CD10 and Ezrin were scarcely detected in the enterocytes from the patient with the MYO5B(G519R) mutation. *MYO5B* and Ras-associated binding (RAB)11A staining overlapped and was found primarily in the subapical region of enterocytes of the MYO5B(G519R) patient. The subapical RAB11A staining pattern seen in this patient differs from previous MVID patients with MYO5B(P660L)⁴ or MYO5B-deficient mouse models,^{7,8} in which RAB11A was diffuse in the cytoplasm. The present MxIF panel is useful for documenting deficits in brush-border transporters in congenital diarrheal diseases using single sections of limited biopsy specimens from infants, and this technique will facilitate an accurate diagnosis and provide novel insights into molecular mechanisms of diseases.

To understand the molecular basis of MVID pathology resulting from

compound heterozygous mutations in *MYO5B* in this patient, we developed a mouse model mimicking the patient's genotype. The 1-step 2-cell embryo microinjection technique⁹ was used to avoid early lethality in homozygous *Myo5b*^{G519R/G519R} mice (Figure 2A and Supplementary Figure 1B). Next, we crossed the *Myo5b*^{G519R/+} mouse with a *Villin-Cre*^{ERT2};*Myo5b*^{fllox/fllox} mouse to generate tamoxifen-inducible, patient-mimicking *VilCre*^{ERT2};*Myo5b*^{G519R/fllox} mice. In this study, the tamoxifen-injected *VilCre*^{ERT2};*Myo5b*^{G519R/fllox} mice are referred to as Myo5b(G519R). After a single tamoxifen injection, Myo5b(G519R) mice showed an average 19% body weight decrease by day 4, similar to that in *VilCre*^{ERT2};*Myo5b*^{fllox/fllox} (*Myo5b* Δ IEC) mice (Figure 2B). The Myo5b(G519R) mouse ileum and colon indicated a severe watery diarrhea phenotype (Figure 2C). Tamoxifen-treated, Cre-lacking *Myo5b*^{G519R/+} mice, which represent the patient's mother's genotype, as well as treated *VilCre*^{ERT2};*Myo5b*^{fllox/+}, displayed healthy intestinal phenotypes (Figure 2B and C) and were used as controls. Myo5b(G519R) intestines showed PAS-positive vesicles, disordered brush-border structures, microvillus inclusions, and subapical accumulation of MYO5B together with Rab11a, recapitulating the MVID patient's phenotype (Figure 2D–G). These observations indicate that the Myo5b(G519R) strain phenocopies the MVID patient.

Important apical sodium transporters, SGLT1, apical sodium-dependent bile acid transporter (ASBT), and NHE3, were internalized away from the apical membrane in Myo5b(G519R) enterocytes (Supplementary Figure 2A–C), consistent with limited absorption of sodium and water, resulting in watery diarrhea. The actinin-4⁺ terminal web structure was disrupted (Supplementary Figure 2F), and the expanded SGLT1 localization with lysosome-associated membrane protein 1 (LAMP1)-expressing lysosomes in the Myo5b(G519R) enterocytes (Supplementary Figure 2A) suggests degradation of mislocalized proteins, similar to the expanded RAB7⁺ vesicles in the MVID patient tissues.




Figure 1. Impaired enterocyte structures and mislocalization of apical proteins in an MVID patient with a point mutation at MYO5B(G519R). (A) Histology of an MVID patient with a MYO5B(G519R) point mutation. Scale bars: 20 μm (upper panels). Transmission electron microscopy (TEM) image of the patient biopsy specimen shows disorganized microvilli, an abnormal accumulation of subapical vesicles, and large lysosomes. Scale bar: 1 μm (lower panel). (B) Immunostaining for the brush-border marker dipeptidyl peptidase 4 (DPP4) (red), a terminal web marker, α -actinin-4 (green), and A-kinase anchoring protein (AKAP)350 (blue) in the patient biopsy specimens. Scale bars: 20 μm . Multiplexed immunofluorescence staining for various epithelial proteins in (C) control pediatric patient biopsy specimens and the (D) MVID patient with a MYO5B(G519R) mutation. MYO5B and RAB11A signals are virtually merged using nuclei signals as a guide. Scale bars: 100 μm in whole biopsy images, 50 μm in cropped images.




Figure 2. Myo5b(G519R) mice recapitulate the MVID patient's phenotype. (A) Genomic editing design to create the G519R mutation in *MYO5B* and breeding strategy for patient-mimicking mouse model using the combination of 1-step 2-cell embryo microinjection and *Cre/loxP* techniques. (B) Significant body weight loss after a tamoxifen injection in *Myo5bΔIEC* and *Myo5b(G519R)* mice. **P* < .0001 vs control by 2-way analysis of variance with Bonferroni multiple comparisons. *n* = 5–6 mice in each group. (C) Diarrhea phenotype in MVID mouse models. No solid feces were observed in the *Myo5b(G519R)* mouse colon. Scale bars: 1 cm. (D) Elongated crypts and blunted villi are present in the *Myo5b(G519R)* mouse duodenum in Alcian Blue/periodic acid–Schiff (PAS) staining. Scale bars: 50 μm; and 10 μm (insets). (E) Phalloidin staining in the duodenum. Confocal volume images (right panels for each mouse) reveal mislocalized microvilli in the inclusions and lateral domain of *Myo5b(G519R)* enterocytes. Scale bars: 50 μm (left) and 10 μm (right). (F) Transmission electron microscopy (TEM) images of crypt and villus cells of the jejunum. In *Myo5b(G519R)* mice, both crypt and villus enterocytes possess blunted and disorganized microvilli and contain numerous sub-apical vesicles. Scale bars: 1 μm. (G) Immunostaining for *MYO5B* and *Rab11a* (sc-166912) in the crypts and villi of small intestine. Scale bars: 10 μm. PAM, roto-spacer-adjacent motif; RNP, ribonucleoprotein; ssDNA, single stranded DNA.

The localization and function of crypt cystic fibrosis transmembrane conductance regulator (CFTR) were intact in Myo5b(G519R) duodenum (Supplementary Figure 2C and D). Üssing chamber experiments showed that CFTR-dependent short-circuit current (I_{sc}), which represents chloride secretion, was up-regulated in steady-state Myo5b(G519R) mouse duodenum. However, bicarbonate secretion, which is likely predominate in villi, showed no significant differences at baseline, after NHE3 inhibition, or forskolin-stimulated conditions (Supplementary Figure 2E).

The Myo5b(G519R) mouse intestine expanded proliferating cell nuclear antigen (PCNA)⁺ crypts and sustained sub-apical olfactomedin 4 (OLFM4) signals in upper crypt and lower villus, suggesting an expansion of immature cells (Supplementary Figure 3A and B). EdU (5-ethynyl-2'-deoxyuridine)⁺ cells reached 90% of the total epithelial height within 24 hours, suggesting more rapid shedding of villus enterocytes in Myo5b(G519R) mice than in healthy controls (Supplementary Figure 3C and D). Indeed, scanning electron microscope images showed immature and disorganized microvilli in Myo5b(G519R) mouse enterocytes (Supplementary Figure 3E). These findings are consistent with a deficit in enterocyte maturation in Myo5b(G519R) mice and in the patient with the MYO5B(G519R) mutation.

In summary, we showed that the compound heterozygous MYO5B(G519R) mutation in an MVID patient and a patient genotype-mimicking, genetically engineered mouse model causes severe diarrhea and disrupts epithelial microvillus structure. This study reveals a methodology for the rapid establishment of a novel MVID mouse model, Myo5b(G519R), to study the effects of compound heterozygous MYO5B mutations *in vivo*, and suggests that the combination of 1-step 2-cell embryo microinjection and Cre/loxP systems is a useful tool for modeling patient-specific monogenic congenital disorders.

ANDREANNA BURMAN*
Department of Cell and Developmental Biology
Vanderbilt University
Nashville, Tennessee, and
Epithelial Biology Center
Vanderbilt University Medical Center
Nashville, Tennessee

MICHAEL MOMOH*
Epithelial Biology Center
Section of Surgical Sciences
Vanderbilt University Medical Center
Nashville, Tennessee

LEESA SAMPSON
JENNIFER SKELTON
Center for Stem Cell Biology
Vanderbilt University
Nashville, Tennessee

JOSEPH T. ROLAND
CYNTHIA RAMOS
Epithelial Biology Center
Section of Surgical Sciences
Vanderbilt University Medical Center
Nashville, Tennessee

EVAN KRYSIOFIK
Department of Cell and Developmental Biology
Vanderbilt University
Nashville, Tennessee

SARI ACRA
Division of Pediatric Gastroenterology
Department of Pediatrics
Vanderbilt University Medical Center
Nashville, Tennessee

JAMES R. GOLDENRING
Department of Cell and Developmental Biology
Vanderbilt University
Nashville, Tennessee, and
Epithelial Biology Center
Section of Surgical Sciences
Vanderbilt University Medical Center
Nashville, Tennessee, and
Nashville VA Medical Center
Nashville, Tennessee

IZUMI KAJI
Department of Cell and Developmental Biology
Vanderbilt University
Nashville, Tennessee, and


Epithelial Biology Center
Section of Surgical Sciences
Vanderbilt University Medical Center
Nashville, Tennessee

References

1. Thiagarajah JR, et al. *Gastroenterology* 2018;154:2045–2059.e6.
2. Aldrian D, et al. *J Clin Med* 2021; 10:481.
3. Bowman DM, et al. *Cell Mol Gastroenterol Hepatol* 2022; 14:553–565.
4. Knowles BC, et al. *J Clin Invest* 2014;124:2947–2962.
5. Kravtsov DV, et al. *Am J Physiol Gastrointest Liver Physiol* 2016; 311:G142–G155.
6. Engevik AC, et al. *Gastroenterology* 2018;155:1883–1897.e10.
7. Schneeberger K, et al. *Proc Natl Acad Sci U S A* 2015;112: 12408–12413.
8. Weis VG, et al. *Cell Mol Gastroenterol Hepatol* 2016;2:131–157.
9. Wu Y, et al. *Nat Commun* 2019; 10:2883.

*These authors contributed equally.

Abbreviations used in this letter: MVID, microvillus inclusion disease; MYO5B, myosin Vb; Myo5b(G519R) mouse, tamoxifen-injected *ViiCre^{ERT2};Myo5b^{G519R/flox}* mouse; MxIF, multiplex immunofluorescence; NHE3, sodium hydrogen antiporter 3; SGLT1, sodium-dependent glucose transporter 1.

 Most current article

© 2023 The Authors. Published by Elsevier Inc. on behalf of the AGA Institute. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).
2352-345X
<https://doi.org/10.1016/j.jcmgh.2022.12.015>

Received October 18, 2022. Accepted December 23, 2022.

Correspondence

Address correspondence to: Izumi Kaji, PhD, Epithelial Biology Center Section of Surgical Sciences, Vanderbilt University Medical Center, 2213 Garland Avenue, MRB IV Room 10465H,

Nashville, Tennessee 37232. e-mail: izumi.kaji@vumc.org.

Acknowledgments

The authors thank Dr. Lane Clarke at the University of Missouri for his expertise in bicarbonate secretion measurements. The authors thank the Vanderbilt Digital Histology (VA Shared Equipment grant 1S1BX003097) and Vanderbilt Cell Imaging Shared Resources (CA68485, DK20593, DK58404, DK59637, and EY08126) for excellent imaging techniques. Generation of Myo5b^{G519R} mice by the Vanderbilt Genome Editing Resource (RRID: SCR_018826) was supported by Cancer

Center Support grant CA68485, the Vanderbilt Diabetes Research and Training Center grant DK020593, and the Center for Stem Cell Biology. Paraffin sectioning and histological staining were performed by the Vanderbilt Translational Pathology Shared Resource (National Cancer Institute/National Institutes of Health Cancer Center Support grant P30 CA68485).

Conflicts of interest

The authors disclose no conflicts.

Funding

This study was supported by National Institutes of

Health (NIH) R01 DK128190, Vanderbilt Digestive Diseases Research Center Pilot and Feasibility grant P30 058404, and the American Physiological Society John F. Perkins, Jr. Research Career Enhancement Award to I.K., and NIH RC2 DK118640, R01 DK48370, and a gift from the Christine Volpe Fund to J.R.G.. A.B. was supported by National Foundation of Science Graduate Research Fellowship Program.

Data Availability Statement

The Myo5b(G519R) mouse strain will be available from the Vanderbilt Cryopreservation Repository.